

List of equations to be used in written examinations in EMP

- Electric field potential and Poisson equation

$$\mathbf{E}(\mathbf{r}) = -\nabla U(\mathbf{r}) \quad \nabla^2 U(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\varepsilon_0} \quad (1)$$

- Electric field of a charge distribution

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\varepsilon_0} \int_{(V)} \frac{\rho(\mathbf{r}')(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} d^3\mathbf{r}' \quad (2)$$

$$U(\mathbf{r}) = \frac{1}{4\pi\varepsilon_0} \int_{(V)} \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d^3\mathbf{r}' \quad (3)$$

- Electric dipole field

$$U(\mathbf{r}) = \frac{\mathbf{p}_e \cdot \mathbf{r}}{4\pi\varepsilon_0 r^3} \quad (4)$$

$$\mathbf{E}(\mathbf{r}) = \frac{3(\mathbf{p}_e \cdot \mathbf{r})\mathbf{r} - \mathbf{p}_e r^2}{4\pi\varepsilon_0 r^5}. \quad (5)$$

$$\mathbf{p}_e = \int \mathbf{r}' \rho(\mathbf{r}') d^3\mathbf{r}' \quad (6)$$

- Electric force on a charge distribution

$$\mathbf{F}_e = \varepsilon_0 \oint_{(\partial V)} \left[\mathbf{E}(\mathbf{E} \cdot \mathbf{n}) - \frac{1}{2} E^2 \mathbf{n} \right] dS \quad (7)$$

- Magnetic force on a current distribution in an external field

$$\mathbf{F}_m = \int_V \mathbf{j}(\mathbf{r}) \times \mathbf{B}(\mathbf{r}) d^3\mathbf{r} \quad (8)$$

- Magnetic field vector potential and Poisson equation

$$\mathbf{B}(\mathbf{r}) = \nabla \times \mathbf{A}(\mathbf{r}) \quad \nabla^2 \mathbf{A}(\mathbf{r}) = -\mu_0 \mathbf{j}(\mathbf{r}) \quad (9)$$

- Biot-Savart law

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{j}(\mathbf{r}') d^3\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} \quad (10)$$

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int_{(V)} \frac{\mathbf{j}(\mathbf{r}') \times (\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} d^3\mathbf{r}' \quad (11)$$

- Inductivity

$$\Phi_i = \sum_k L_{ik} I_k \quad (12)$$

- Mutual inductivity of two current loops

$$L_{ik} = \frac{\mu_0}{4\pi} \oint_{i,k} \frac{d\mathbf{l}_i d\mathbf{l}_k}{|\mathbf{r}(l_i) - \mathbf{r}(l_k)|} \quad (13)$$

- Magnetic dipole field

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \frac{\mathbf{p}_m \times \mathbf{r}}{r^3} \quad (14)$$

$$\mathbf{B} = \frac{\mu_0}{4\pi} \frac{3\mathbf{r}(\mathbf{p}_m \cdot \mathbf{r}) - \mathbf{p}_m r^2}{r^5} \quad (15)$$

$$\mathbf{p}_m = \frac{1}{2} \int \mathbf{r}' \times \mathbf{j}(\mathbf{r}') d^3 \mathbf{r}' \quad (16)$$

- Magnetic force on a current distribution

$$\mathbf{F}_m = \frac{1}{\mu_0} \oint_{(\partial V)} \left[\mathbf{B}(\mathbf{B} \cdot \mathbf{n}) - \frac{1}{2} B^2 \mathbf{n} \right] dS \quad (17)$$

- Electric conductivity

$$\mathbf{j} = \sigma \mathbf{E} \quad (18)$$

- Electric current in a closed loop

$$\dot{U} = L\ddot{I} + R\dot{I} + \frac{I}{C} \quad (19)$$

- Maxwell equations in vacuum

$$\begin{aligned} \nabla \cdot \mathbf{E} &= \frac{\rho}{\epsilon_0} & \nabla \cdot \mathbf{B} &= 0 \\ \nabla \times \mathbf{E} &= -\frac{\partial \mathbf{B}}{\partial t} & \nabla \times \mathbf{B} &= \mu_0 \mathbf{j} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \end{aligned} \quad (20)$$

- Continuity equation

$$\nabla \cdot \mathbf{j} = -\frac{\partial \rho}{\partial t} \quad (21)$$

- Poynting theorem

$$\frac{\partial w}{\partial t} + \nabla \cdot \mathbf{P} + \mathbf{j} \cdot \mathbf{E} = 0 \quad (22)$$

$$\mathbf{P} = \frac{1}{\mu_0} (\mathbf{E} \times \mathbf{B}) \quad w = \frac{1}{2} \epsilon_0 E^2 + \frac{1}{2} \frac{B^2}{\mu_0} \quad (23)$$

- Electromagnetic field momentum and angular momentum densities

$$\mathbf{g} = \epsilon_0 (\mathbf{E} \times \mathbf{B}) \quad \gamma = \mathbf{r} \times \mathbf{g} \quad (24)$$

- Maxwell equations in matter

$$\begin{aligned} \nabla \cdot \mathbf{D} &= \rho & \nabla \cdot \mathbf{B} &= 0 \\ \nabla \times \mathbf{E} &= -\frac{\partial \mathbf{B}}{\partial t} & \nabla \times \mathbf{H} &= \mathbf{j} + \frac{\partial \mathbf{D}}{\partial t} \end{aligned} \quad (25)$$

- Electric fields in matter

$$\rho_v = -\nabla \cdot \mathbf{P} \quad \sigma_v = \mathbf{P} \cdot \mathbf{n} \quad (26)$$

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P} \quad \mathbf{D} = \epsilon_0 \epsilon \mathbf{E} \quad \mathbf{P} = \epsilon_0 (\epsilon - 1) \mathbf{E} \quad (27)$$

- Magnetic fields in matter

$$\mathbf{H} = \frac{\mathbf{B}}{\mu_0} - \mathbf{M} \quad \mathbf{H} = \frac{\mathbf{B}}{\mu \mu_0} \quad \mathbf{M} = (1 - \frac{1}{\mu}) \frac{\mathbf{B}}{\mu_0} \quad (28)$$

- Electromagnetic wave in vacuum

$$\mathbf{E}(\mathbf{r},t) = \mathbf{E}_0 e^{i(kz-\omega t)} \quad c_0 \mathbf{B}(\mathbf{r},t) = \hat{\mathbf{e}}_z \times \mathbf{E}(\mathbf{r},t) \quad \omega = kc_0 \quad (29)$$

- Electromagnetic wave in a waveguide

$$\nabla^2 \mathbf{E}(\mathbf{r},t) - \frac{1}{c^2} \frac{\partial^2 \mathbf{E}(\mathbf{r},t)}{\partial t^2} = 0 \quad \nabla^2 \mathbf{H}(\mathbf{r},t) - \frac{1}{c^2} \frac{\partial^2 \mathbf{H}(\mathbf{r},t)}{\partial t^2} = 0 \quad (30)$$

$$\mathbf{E}(\mathbf{r},t) = \mathbf{E}(\rho) e^{i(kz-\omega t)} \quad \mathbf{H}(\mathbf{r},t) = \mathbf{H}(\rho) e^{i(kz-\omega t)} \quad (31)$$

$$\left(\nabla_{\perp}^2 + \frac{\omega^2}{c^2} - k^2 \right) \mathbf{E}(\rho) = 0 \quad \left(\nabla_{\perp}^2 + \frac{\omega^2}{c^2} - k^2 \right) \mathbf{H}(\rho) = 0 \quad (32)$$

- TEM wave in a waveguide

$$\omega = kc \quad \mathbf{H}(\rho) = Z^{-1} \hat{\mathbf{e}}_z \times \mathbf{E}(\rho) \quad Z = \sqrt{\frac{\mu_0}{\epsilon \epsilon_0}} \quad \nabla_{\perp}^2 \mathbf{E}(\rho) = 0 \quad \nabla_{\perp}^2 \mathbf{H}(\rho) = 0 \quad (33)$$

- Radiation field of a current distribution

$$\begin{aligned} \mathbf{B}(\mathbf{r},t) &= -\frac{\mu_0}{4\pi c_0 r} \frac{\mathbf{r}}{r} \times \int_{V'} \frac{\partial \mathbf{j}(\mathbf{r}', t - \frac{r}{c_0} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r c_0})}{\partial t} d^3 \mathbf{r}' \\ \mathbf{E}(\mathbf{r},t) &= -\frac{\mathbf{r}}{r} \times c_0 \mathbf{B}(\mathbf{r},t) \end{aligned} \quad (34)$$

- Dipole radiation of an electric and magnetic dipole

$$\begin{aligned} \mathbf{B}(\mathbf{r},t) &= -\frac{\mu_0}{4\pi c_0 r} \frac{\mathbf{r}}{r} \times \ddot{\mathbf{p}}_e(t - \frac{r}{c_0}) & \mathbf{E}(\mathbf{r},t) &= -\frac{\mathbf{r}}{r} \times c_0 \mathbf{B}(\mathbf{r},t) \\ \mathbf{B}(\mathbf{r},t) &= -\frac{\mu_0}{4\pi c_0^2 r} \frac{\mathbf{r}}{r} \times \left(\frac{\mathbf{r}}{r} \times \ddot{\mathbf{p}}_m(t - \frac{r}{c_0}) \right) & \mathbf{E}(\mathbf{r},t) &= -\frac{\mathbf{r}}{r} \times c_0 \mathbf{B}(\mathbf{r},t) \end{aligned} \quad (35)$$